




Unexpected Features of Solar Gamma-Ray Emission














A Completely Data Driven Background Model



$$N_b(T_x, T_y) = \sum_{\text{RA,Dec}} \frac{N_o(\text{RA,Dec})}{\epsilon(\text{RA,Dec})} \times \sum_{\text{t}} \epsilon^*(\text{RA,Dec}, T_x, T_y, t)$$





How are solar gamma-rays produced?




Solar Flare gamma-rays are low energy ($E_{max} = 4 \text{ GeV}$)









First Observations of Solar Halo Gamma Rays Over a Full Solar Cycle

Tim Linden (1,1,2,* Jung-Tsung Li (3,3 Bei Zhou (3,4,5 Isabelle John (3,6,7 Milena Crnogorčević (3,1 Annika H. G. Peter (3,3,8,9 and John F. Beacom (3,8,9))


Stockholm University and The Oskar Klein Centre for Cosmoparticle Physics, Alba Nova, 10691 Stockholm, Sweden Erlangen Centre for Astroparticle Physics (ECAP), Friedrich-Alexander-Universität Erlangen-Nürnberg, Nikolaus-Fiebiger-Str. 2, 91058 Erlangen, Germany


Center for Cosmology and AstroParticle Physics, The Ohio State University, Columbus, OH 43210, USA Theory Division, Fermi National Accelerator Laboratory, Batavia, IL 60510, USA Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA Dipartimento di Fisica, Università degli Studi di Torino, via P. Giuria, 1 10125 Torino, Italy Pepartment of Physics, The Ohio State University, Columbus, OH 43210, USA Department of Astronomy, The Ohio State University, Columbus, OH 43210, USA


We analyze 15 years of *Fermi*-LAT data and produce a detailed model of the Sun's inverse-Compton scattering emission (solar halo), which is powered by interactions between ambient cosmic-ray electrons and positrons with sunlight. By developing a novel analysis method to analyze moving sources, we robustly detect the solar halo at energies between 31.6 MeV and 100 GeV, and angular extensions up to 45° from the Sun, providing new insight into spatial regions where there are no direct measurements of the galactic cosmic-ray flux. The large statistical significance of our signal allows us to sub-divide the data and provide the first γ -ray probes into the time-variation and azimuthal asymmetry of the solar modulation potential, finding time-dependent changes in solar modulation both parallel and perpendicular to the ecliptic plane. Our results are consistent with (but with independent uncertainties from) local cosmic-ray measurements, unlocking new probes into both astrophysical and beyond-standard-model processes near the solar surface.


Extremely well measured inverse-Compton spectrum and morphology.


30 MeV-100 GeV in energy 0-45 degrees in radius




















First Observation of Time Variation in the Solar-Disk Gamma-Ray Flux with Fermi

Kenny C. Y. Ng,^{1,2,*} John F. Beacom,^{1,2,3,†} Annika H. G. Peter,^{1,2,3,‡} and Carsten Rott^{4,§}

¹Center for Cosmology and AstroParticle Physics (CCAPP), Ohio State University, Columbus, OH 43210

²Department of Physics, Ohio State University, Columbus, OH 43210

³Department of Astronomy, Ohio State University, Columbus, OH 43210

⁴Department of Physics, Sungkyunkwan University, Suwon 440-746, Korea

(Dated: 7 November 2015)

The solar disk is a bright gamma-ray source. Surprisingly, its flux is about one order of magnitude higher than predicted. As a first step toward understanding the physical origin of this discrepancy, we perform a new analysis in 1–100 GeV using 6 years of public Fermi-LAT data. Compared to the previous analysis by the Fermi Collaboration, who analyzed 1.5 years of data and detected the solar disk in 0.1–10 GeV, we find two new and significant results: 1. In the 1–10 GeV flux (detected at $> 5\sigma$), we discover a significant time variation that anticorrelates with solar activity. 2. We detect gamma rays in 10–30 GeV at $> 5\sigma$, and in 30–100 GeV at $> 2\sigma$. The time variation strongly indicates that solar-disk gamma rays are induced by cosmic rays and that solar atmospheric magnetic fields play an important role. Our results provide essential clues for understanding the underlying gamma-ray production processes, which may allow new probes of solar atmospheric magnetic fields, cosmic rays in the solar system, and possible new physics. Finally, we show that the Sun is a promising new target for ground-based TeV gamma-ray telescopes such as HAWC and LHAASO.

PACS numbers: 95.85.Pw, 96.50.S-, 13.85.Qk, 96.50.Vg

I. INTRODUCTION

The Sun is well studied and understood with a broad set of messengers at different energies. For example, the optical photon and MeV neutrino spectra confirm a de-

angular size of the Sun ($\simeq 0.5^{\circ}$); we denote it (plus any potential non-cosmic-ray contribution) as the solar-disk component.

Theoretical estimation of both components requires taking into account the effects of solar magnetic activity.

Unexpected Dip in the Solar Gamma-Ray Spectrum

Qing-Wen Tang,^{1,2,*} Kenny C. Y. Ng,^{3,†} Tim Linden,^{1,‡}
Bei Zhou,^{1,4,§} John F. Beacom,^{1,4,5,¶} and Annika H. G. Peter^{1,4,5,**}

¹ Center for Cosmology and AstroParticle Physics (CCAPP),

Ohio State University, Columbus, Ohio 43210, USA

² Department of Physics, Nanchang University, Nanchang 330031, China

³ Department of Particle Physics and Astrophysics,

Weizmann Institute of Science, Rehovot 76100, Israel

⁴ Department of Physics, Ohio State University, Columbus, Ohio 43210, USA

⁵ Department of Astronomy, Ohio State University, Columbus, Ohio 43210, USA

(Dated: 9th May 2018)

The solar disk is a bright source of multi-GeV gamma rays, due to the interactions of hadronic cosmic rays with the solar atmosphere. However, the underlying production mechanism is not understood, except that its efficiency must be greatly enhanced by magnetic fields that redirect some cosmic rays from ingoing to outgoing before they interact. To elucidate the nature of this emission, we perform a new analysis of solar atmospheric gamma rays with 9 years of Fermi-LAT data, which spans nearly the full 11-year solar cycle. We detect significant gamma-ray emission from the solar disk from 1 GeV up to $\gtrsim 200\,\mathrm{GeV}$. The overall gamma-ray spectrum is much harder ($\sim E_{\gamma}^{-2.2}$) than the cosmic-ray spectrum ($\sim E_{\mathrm{CR}}^{-2.7}$). We find a clear anticorrelation between the solar cycle phase and the gamma-ray flux between 1–10 GeV. Surprisingly, we observe a spectral dip between $\sim 30-50\,\mathrm{GeV}$ in an otherwise power-law spectrum. This was not predicted, is not understood, and may provide crucial clues to the gamma-ray emission mechanism. The flux above 100 GeV, which is brightest during the solar minimum, poses exciting opportunities for HAWC, LHAASO, IceCube, and KM3NeT.

I. INTRODUCTION

particles accelerated during energetic solar events, such as solar flares and coronal mass ejections [11], can pro-

Evidence for a New Component of High-Energy Solar Gamma-Ray Production

Tim Linden,^{1,*} Bei Zhou,^{1,2,†} John F. Beacom,^{1,2,3,‡} Annika H. G. Peter,^{1,2,3,§} Kenny C. Y. Ng,^{4,¶} and Qing-Wen Tang^{1,5,**}

¹Center for Cosmology and AstroParticle Physics (CCAPP), The Ohio State University, Columbus, OH 43210

²Department of Physics, The Ohio State University, Columbus, OH 43210

³Department of Astronomy, The Ohio State University, Columbus, OH 43210

⁴Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 76100, Israel

⁵Department of Physics, Nanchang University, Nanchang 330031, China

The observed multi-GeV gamma-ray emission from the solar disk — sourced by hadronic cosmic rays interacting with gas, and affected by complex magnetic fields — is not understood. Utilizing an improved analysis of the Fermi-LAT data that includes the first resolved imaging of the disk, we find strong evidence that this emission is produced by two separate mechanisms. Between 2010–2017 (the rise to and fall from solar maximum), the gamma-ray emission is dominated by a polar component. Between 2008–2009 (solar minimum) this component remains present, but the total emission is instead dominated by a new equatorial component with a brighter flux and harder spectrum. Most strikingly, although 6 gamma rays above 100 GeV are observed during the 1.4 years of solar minimum, none are observed during the next 7.8 years. These features, along with a 30–50 GeV spectral dip which will be discussed in a companion paper, were not anticipated by theory. To understand the underlying physics, Fermi and HAWC observations of the imminent Cycle 25 solar minimum are crucial.

The Sun is a bright source of multi-GeV γ -rays, with emission observed both from its halo — due to cosmic-rays electrons interacting with solar photons — and its disk — due to hadronic cosmic rays (mostly protons) interacting with solar gas. (Emission from solar particle acceleration is only bright during flares and has not been observed above 4 GeV [1–8].) Although the halo emission [9] agrees with theory [10–12], the disk emission does not, and hence is our focus.

Until recently the most extensive analysis of solar disk

is detected up to ~ 30 GeV. Most significantly, we discover a spectral dip between 30–50 GeV. This dip is unexpected and its origin is unknown. Here we extend the analyses of Refs. [13, 17] by going to higher energies, studying the time variation in a new way, and performing the first analysis of flux variations across the resolved solar disk. In the following, we detail our methodology, highlight key discoveries, and discuss their possible theoretical implications.

The importance of this work is manifold. Decouse the disk

First Observations of Solar Disk Gamma Rays over a Full Solar Cycle

Tim Linden,^{1,2,*} John F. Beacom,^{2,3,4,†} Annika H. G. Peter,^{2,3,4,‡} Benjamin J. Buckman,^{2,3,§} Bei Zhou,^{2,3,5,¶} and Guanying Zhu^{2,3,**}


¹Stockholm University and The Oskar Klein Centre for Cosmoparticle Physics, AlbaNova, 10691 Stockholm, Sweden ²Center for Cosmology and AstroParticle Physics (CCAPP), Ohio State University, Columbus, Ohio 43210, USA ³Department of Physics, Ohio State University, Columbus, Ohio 43210, USA ⁴Department of Astronomy, Ohio State University, Columbus, Ohio 43210, USA ⁵Department of Physics and Astronomy, Johns Hopkins University, Baltimore, Maryland 21218, USA


The solar disk is among the brightest γ -ray sources in the sky. It is also among the most mysterious. No existing model fully explains the luminosity, spectrum, time variability, and morphology of its emission. We perform the first analysis of solar-disk γ -rays over a full 11-year solar cycle, utilizing a powerful new method to differentiate solar signals from astrophysical backgrounds. We produce: (i) a robustly measured spectrum from 100 MeV to 100 GeV, reaching a precision of several percent in the 1–10 GeV range, (ii) new results on the anti-correlation between solar activity and γ -ray emission, (iii) strong constraints on short-timescale variability, ranging from hours to years, and (iv) new detections of the equatorial and polar morphologies of high-energy γ -rays. Intriguingly, we find no significant energy dependence in the time variability of solar-disk emission, indicating that strong magnetic-field effects close to the solar surface, rather than modulation throughout the heliosphere, must primarily control the flux and morphology of solar-disk emission.

I. INTRODUCTION

The Sun is a special astrophysical source. Its close proximity allows detailed studies critical to understanding other stars. The ability to spatially resolve solar emission is especially important for probing high-energy, nonthermal processes, which can be highly local. These processes reveal charged-particle


The Sun's γ -ray emission is dramatically affected by its magnetic fields. Without magnetic fields, the disk emission would have two components. At energies above ~ 1 GeV, the γ -ray direction increasingly follows that of the parent cosmic ray. Accordingly, only cosmic rays that graze the solar surface can interact and have the γ -rays escape [14]. The corresponding emission from the solar limb is too faint to be observed






Intensity







Intensity Spectrum



Gamma-Rays Continue up to 3 TeV during Solar Minimum!



Low-Energy Spectrum does not cut off either!




Intensity

Spectrum

Time Variability

X




Intensity


Spectrum


Time Variability

Morphology



Intensity
Spectrum
Time Variability
Morphology





Intensity

Spectrum

X

Time Variability

Morphology

X


Spectral Dip

X


So basically everything is wrong....

How do we model this?


$$\Phi_{\odot}(E_{\gamma}) = \pi R_{\odot}^2 \Phi_{\rm CR}(E_{\rm CR}) C(E_{\gamma}, E_{\rm CR}) f_{\rm sur} f_{\rm turn} f_{\rm int}$$




$$\Phi_{\odot}(E_{\gamma}) = \pi R_{\odot}^2 \Phi_{\rm CR}(E_{\rm CR}) C(E_{\gamma}, E_{\rm CR}) f_{\rm sur} f_{\rm turn} f_{\rm int}$$



$$\Phi_{\odot}(E_{\gamma}) = \pi R_{\odot}^2 \Phi_{\rm CR}(E_{\rm CR}) C(E_{\gamma}, E_{\rm CR}) f_{\rm sur} f_{\rm turn} f_{\rm int}$$



$$\Phi_{\odot}(E_{\gamma}) = \pi R_{\odot}^2 \Phi_{\rm CR}(E_{\rm CR}) C(E_{\gamma}, E_{\rm CR}) f_{\rm sur} f_{\rm turn} f_{\rm int}$$



We see, but we don't understand.

Help?

Solar disk gamma-rays provide a new handle into fundamental questions in solar magnetohydrodynamics.

Huge Potential For Cross-Correlations with other wavelengths

