


# THE SEARCH FOR DARK MATTER ANNIHILATION IN THE GAMMA-RAY SKY

TIM LINDEN


WITH: TANSU DAYLAN, DOUG FINKBEINER, DAN HOOPER, STEPHEN PORTILLO, NICK RODD, TRACY SLATYER, ILIAS CHOLIS, MANOJ KAPLINGHAT, HAIBO YU, PHILIPP MERTSCH AND OTHERS

# DARK MATTER




(b)

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison-Wesley.



# DARK MATTER

WEAKLY INTERACTING MASSIVE PARTICLES



Ignoring several possible complications, a particle with a weak interaction cross-section and a mass on the weak scale is expected to naturally obtain the correct relic abundance through thermal freeze-out in the early universe



$$\left( \frac{\Omega_\chi}{0.2} \right) \simeq \frac{x_{f.o.}}{20} \left( \frac{10^{-8} \text{ GeV}^{-2}}{\sigma} \right)$$

$$\langle \sigma v \rangle \sim 10^{-8} \text{ GeV}^{-2} (3 \times 10^{-28} \text{ GeV}^2 \text{ cm}^2) 10^{10} \frac{\text{cm}}{\text{s}} = 3 \times 10^{-26} \frac{\text{cm}^3}{\text{s}}$$

# DARK MATTER

WEAKLY INTERACTING MASSIVE PARTICLES

INDIRECT DETECTION OF WIMPS



# INDIRECT DETECTION OF WIMPS


**Astrophysics**

**Particle Physics**




**Instrumental Response**

# INDIRECT DETECTION OF WIMPS

 **WE HAVE AN INSTRUMENT!**

Launched in June 2008 and has been taking science data for > 6 yr

Detects  $\gamma$ -rays between  $\sim 30$  MeV - 1 TeV



Effective Area  $\sim 0.8$  m<sup>2</sup>

Field of View  $\sim 2.4$  sr

Energy Resolution  $\sim 10\%$


Angular Resolution is highly Energy Dependent

# INDIRECT DETECTION OF WIMPS

→ PARTICLE PHYSICS

## Why Do We Search in Gamma-Rays?

For a dark matter particle with a mass of  $\sim 100$  GeV, the standard model annihilation products tend to have energy in the 10 GeV range



# INDIRECT DETECTION OF WIMPS



## DARK MATTER DENSITY PROFILES

$$\rho_{\text{NFW}} = \left( \frac{r}{r_s} \right)^{-\gamma} \left( 1 + \frac{r}{r_s} \right)^{-3+\gamma}$$

A simple analytic formula has been found that provides a reasonable fit to the observed density distribution of dark matter over halos of widely varying masses.

In the standard NFW scenario,  $\gamma = 1$

Navarro, Frenk, White (1996)


Springel et al. (2008, 0809.0898)



# INDIRECT DETECTION OF WIMPS

→ DARK MATTER DENSITY PROFILES

→ THE GALACTIC CENTER



For typical parameters from an  
NFW profile:

$$J \sim 10^{21} \text{ GeV}^2 \text{ cm}^{-5}$$

Name	GLON (deg)	GLAT (deg)	Distance (kpc)	$\overline{\log_{10}(J_{\text{NFW}})^a}$ ( $\log_{10}[\text{GeV}^2 \text{ cm}^{-5} \text{ sr}]$ )
Bootes I	358.1	69.6	66	$18.8 \pm 0.22$
Bootes II	353.7	68.9	42	–
Bootes III	35.4	75.4	47	–
Canes Venatici I	74.3	79.8	218	$17.7 \pm 0.26$
Canes Venatici II	113.6	82.7	160	$17.9 \pm 0.25$
Canis Major	240.0	-8.0	7	–
Carina	260.1	-22.2	105	$18.1 \pm 0.23$
Coma Berenices	241.9	83.6	44	$19.0 \pm 0.25$
Draco	86.4	34.7	76	$18.8 \pm 0.16$
Fornax	237.1	-65.7	147	$18.2 \pm 0.21$
Hercules	28.7	36.9	132	$18.1 \pm 0.25$
Leo I	226.0	49.1	254	$17.7 \pm 0.18$
Leo II	220.2	67.2	233	$17.6 \pm 0.18$
Leo IV	265.4	56.5	154	$17.9 \pm 0.28$
Leo V	261.9	58.5	178	–
Pisces II	79.2	-47.1	182	–
Sagittarius	5.6	-14.2	26	–
Sculptor	287.5	-83.2	86	$18.6 \pm 0.18$
Segue 1	220.5	50.4	23	$19.5 \pm 0.29$
Segue 2	149.4	-38.1	35	–
Sextans	243.5	42.3	86	$18.4 \pm 0.27$
Ursa Major I	159.4	54.4	97	$18.3 \pm 0.24$
Ursa Major II	152.5	37.4	32	$19.3 \pm 0.28$
Ursa Minor	105.0	44.8	76	$18.8 \pm 0.19$
Willman 1	158.6	56.8	38	$19.1 \pm 0.31$

# INDIRECT DETECTION OF WIMPS

→ DARK MATTER DENSITY PROFILES

Put Another Way:


→ THE GALACTIC CENTER

The Fermi-LAT telescope observes a flux in the inner  $1^\circ$  between 1-3 GeV of approximately  $1 \times 10^{-10}$  erg cm<sup>-2</sup> s<sup>-1</sup>

A Generic Dark Matter Scenario predicts a flux of  $2 \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup>

Unfortunately, the backgrounds are not negligible.

Chandra image of Galactic Center



# INDIRECT DETECTION OF WIMPS



PUTTING IT ALL TOGETHER

$$\phi_s(\Delta\Omega) = \underbrace{\frac{1}{4\pi} \frac{\langle\sigma v\rangle}{2m_{\text{DM}}^2} \int_{E_{\text{min}}}^{E_{\text{max}}} \frac{dN_\gamma}{dE_\gamma} dE_\gamma}_{\Phi_{\text{PP}}} \times \underbrace{\int_{\Delta\Omega} \left\{ \int_{\text{l.o.s.}} \rho^2(\mathbf{r}) dl \right\} d\Omega'}_{\text{J-factor}}$$

Fortunately, these terms are separable for standard CDM

# **WHY WE'RE DOING WHAT WE'RE DOING**

- 1.) Dark Matter is one of the most mysterious, but important extensions to the standard model**
- 2.) WIMPs are among the most well-motivated models for dark matter annihilation**
- 3.) The observation of dark matter annihilation products offers the capability to observe/understand the WIMP particle**
- 4.) The Milky Way Galactic Center is a promising target for indirect detection studies.**
- 5.) The Fermi-LAT has provided us with an unparalleled ability to detect WIMPs annihilating at the thermal cross-section**

# PREVIOUS WORK


Many Analyses of the Galactic Center over the past 5 years:

Goodenough & Hooper (2009)	0910.2998
Hooper & Goodenough (2011, PLB 697 412)	1010.2752
Hooper & TL (2011, PRD 84 12)	1110.0006
Abazajian & Kaplinghat (2012, PRD 86 8)	1207.6047
Hooper & Slatyer (2013, PDU 2 18)	1302.6589
Gordon & Macias (2013, PRD 8 8)	1306.5725
Macias & Gordon (2013, PRD 89 6)	1312.6671
Abazajian et al. (2014, PRD 90 2)	1402.4090
Daylan et al. (2014)	1402.6703
Calore et al. (2014)	1409.0042

Different studies have used various techniques and regions of interest, but have obtained consistent results!

# EXCESS SPECTRUM



Calore et al. (2014, 1409.0042)



**Spectral Model highly resilient to changing systematic background models ~300 models considered here.**

**Low energy spectrum hard to constrain due to systematics**  
**High energy spectrum difficult due to statistics**

# MORPHOLOGY




Inner galaxy prefers density profile  $\gamma = 1.26$

Galactic Center prefers  $\gamma = 1.17$

$\gamma = 1.26$  is consistent with both, profile may also vary with radius


# MORPHOLOGY



Can additionally fix the spectrum and allow the normalization to float independently in different radial bins. In this case we find  $\gamma = 1.4$ , which provides some evidence that the profile is steepening with distance.




# MORPHOLOGY




Calore et al (2014) also find evidence for excess emission in morphological bins that extend far from the GC.


Consistent spectrum!



# EMISSION MAPS




INNER GALAXY




GALACTIC CENTER

# PROFILE CORES



**The emission intensity continues to rise to within 10 pc of the GC.**



# CENTERED-NESS?



The center of the emission profile is located to within  $0.05^\circ$  of Sgr A\*.

This disfavors Sgr A East as the source of the  $\gamma$ -ray excess (though only at  $2\sigma$ ).

# ELLIPTICITY



The ellipticity serves as a powerful discriminator of baryonic mechanisms, which tend to be much more luminous along the plane.

# CURRENT STATE OF MEASUREMENTS

All published studies agree:

- The spectrum of the excess is peaked at an energy of  $\sim 2$  GeV, and falls off at low energies with a spectrum that is harder than expected for astrophysical pion emission
- The excess extends to at least  $10^\circ$  away from the galactic center, following a 3D profile which falls in intensity as  $r^{-2.2}$  to  $-2.8$

# IMPORTANT CAVEAT

I have discussed “dark matter fits” to the  $\gamma$ -ray data.

But this does NOT mean that the mechanism producing the excess has a dark matter origin


The data analysis tells us that the model of  $\gamma$ -ray data improves when we add a template with:

- A spherically symmetric, radially falling emission profile with  $r^{-2.0}$  to  $-2.8$
- A spectrum which peaks at an energy of  $\sim 2$  GeV and has a hard low-energy spectrum compared to known astrophysical emission mechanisms

# INTERPRETATIONS

**Three Classes of Interpretations Have Been Proposed So Far:**


**1.) A Population of GC Millisecond Pulsars**



**2.) An Outburst of Hadronic or Leptonic Emission from the Galactic Center**



**3.) Dark Matter Annihilation**






# INTERPRETATIONS

## MILLISECOND PULSARS

- To first order, the peak of the MSP energy spectrum matches the peak of the observed excess
- MSPs are thought to be overabundant in dense star-forming regions (like globular clusters, and potentially the galactic center)



ABAZAJIAN (2011, 1011.4275)


ABAZAJIAN & KAPLINGHAT (2012, 1207.6047)

PETROVIC ET AL. (2014, 1411.2980)

# INTERPRETATIONS

## MILLISECOND PULSARS: M31

### DEGENERACY WITH MILLI-SECOND PULSARS IN SPATIAL PROFILE



We make the reasonable assumption that Low-Mass X-ray Binaries have the same spatial distribution as MSPs


400'' towards M31 center =  
1.5 kpc distance from center =  
10 degrees towards MW center

Orange line is same as best-fit excess template ( $R^{-1.2}$  in projection implies  $r^{-2.2}$  de-projected)!

Slide from Manoj Kaplinghat

# INTERPRETATIONS

## MILLISECOND PULSARS




- Analyze the average spectrum and luminosity of the Fermi MSP and globular cluster populations:
  - 5.5 years of data
  - P7 Reprocessed Photons
  - 15 energy bins, no spectral model assumed

CHOLIS, TL, HOOPER (2014, 1407.5583)  
CHOLIS, TL, HOOPER (2014, 1407.5625)

# INTERPRETATIONS

MILLISECOND PULSARS




CHOLIS, TL, HOOPER (2014, 1407.5583)  
CHOLIS, TL, HOOPER (2014, 1407.5625)

# INTERPRETATIONS

## MILLISECOND PULSARS


- There would need to be 226 (+91/-67) MSPs with luminosity  $> 10^{34}$  erg s $^{-1}$  in the circular region, and 61.9 (+60/-33.7) with luminosity  $> 10^{35}$  erg s $^{-1}$ .
- These should be detectable by the Fermi-LAT as bright point sources
- We can also compare the MSP population to observed LMXBs. The ratio of LMXBs to the MSP luminosity of globular clusters, predicts a population of 103 (+70/-45) LMXBs in the GC in order to produce the GC excess. Only 6 are observed.



# INTERPRETATIONS


## MILLISECOND PULSARS

- Petrovic et al. argue that this may still be consistent with the data, if a break in the MSP luminosity function is added in order to decrease the number of bright systems.
- It is not clear how this new cutoff is affected by non-isotropic emission “beaming”, which is expected to exist in most pulsars.



# INTERPRETATIONS

→ HADRONIC OUTBURSTS



CARLSON & PROFUMO (2014, 1405.7685)

## Echoes of multiple outbursts of Sagittarius A\* revealed by Chandra

M. Clavel<sup>1,2</sup>, R. Terrier<sup>1</sup>, A. Goldwurm<sup>1,2</sup>, M. R. Morris<sup>3</sup>, G. Ponti<sup>4</sup>, S. Soldi<sup>1</sup>, and G. Trap<sup>5,1</sup>

<sup>1</sup> AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/DSM, Observatoire de Paris, Paris Cité ; 10, rue Alice Domon et Léonie Duquet, 75205 Paris Cedex 13, France  
<sup>2</sup> Centre d'Astrophysique et Astronomie, Université of California, Los Angeles, CA 90095-1547, USA  
<sup>3</sup> Lawrence Livermore National Laboratory, Livermore, CA 94550, USA  
<sup>4</sup> Institut für Experimentelle und Angewandte Physik, Universität Wien, A-1040 Vienna, Austria  
<sup>5</sup> Institut für Experimentelle und Angewandte Physik, Universität Wien, A-1040 Vienna, Austria

Accepted 12 July 2013

### ABSTRACT





and temporal variability of the X-ray radiation from the central supermassive black hole, Sagittarius A\*, is studied. This emission on small angular scales is observed from 1999 to 2011. Towards positive latitudes, the emission at 6.4 keV is observed.

# INTERPRETATIONS



## HADRONIC OUTBURSTS


Difficult to explain the low-energy spectrum without introducing highly peaked proton injection spectra





# INTERPRETATIONS

→ HADRONIC OUTBURSTS



Best Fitting Linear Combination of Hadronic Outburst Models:


TS=51 (14 d.o.f)

Best Fitting NFW Template

TS=315 (5 d.o.f)

# INTERPRETATIONS

## LEPTONIC OUTBURSTS



Electron Cooling is a significant issue — the models which correctly fit the morphology of the GC excess are poor fits to the spectrum of the GC excess, and vice versa.

# INTERPRETATIONS







## ASTROPHYSICAL MECHANISMS: BAYESIAN VIEW

- **Current astrophysical models form a relatively poor fit to the excess.**
- **However, the Bayesian prior on the existence of these emission mechanisms is quite high.**
- **Preview - More nuanced astrophysical models should be coming soon, and may provide reasonable fits to the data. Stay tuned!**

# INTERPRETATIONS

 **DARK MATTER**





# INTERPRETATIONS



## DARK MATTER

BERLIN, HOOPER, MCDERMOTT (2014)



Model Number	DM	Mediator	Interactions	Elastic Scattering	Near Future Reach?	
					Direct	LHC
1	Dirac Fermion	Spin-0	$\bar{\chi}\gamma^5\chi, \bar{f}f$	$\sigma_{SI} \sim (q/2m_\chi)^2$ (scalar)	No	Maybe
1	Majorana Fermion	Spin-0	$\bar{\chi}\gamma^5\chi, \bar{f}f$	$\sigma_{SI} \sim (q/2m_\chi)^2$ (scalar)	No	Maybe
2	Dirac Fermion	Spin-0	$\bar{\chi}\gamma^5\chi, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q^2/4m_n m_\chi)^2$	Never	Maybe
2	Majorana Fermion	Spin-0	$\bar{\chi}\gamma^5\chi, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q^2/4m_n m_\chi)^2$	Never	Maybe
3	Dirac Fermion	Spin-1	$\bar{\chi}\gamma^\mu\chi, \bar{b}\gamma_\mu b$	$\sigma_{SI} \sim \text{loop (vector)}$	Yes	Maybe
4	Dirac Fermion	Spin-1	$\bar{\chi}\gamma^\mu\chi, \bar{f}\gamma_\mu\gamma^5f$	$\sigma_{SD} \sim (q/2m_n)^2$ or $\sigma_{SD} \sim (q/2m_\chi)^2$	Never	Maybe
5	Dirac Fermion	Spin-1	$\bar{\chi}\gamma^\mu\gamma^5\chi, \bar{f}\gamma_\mu\gamma^5f$	$\sigma_{SD} \sim 1$	Yes	Maybe
5	Majorana Fermion	Spin-1	$\bar{\chi}\gamma^\mu\gamma^5\chi, \bar{f}\gamma_\mu\gamma^5f$	$\sigma_{SD} \sim 1$	Yes	Maybe
6	Complex Scalar	Spin-0	$\phi^\dagger\phi, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q/2m_n)^2$	No	Maybe
6	Real Scalar	Spin-0	$\phi^2, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q/2m_n)^2$	No	Maybe
6	Complex Vector	Spin-0	$B_\mu^\dagger B^\mu, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q/2m_n)^2$	No	Maybe
6	Real Vector	Spin-0	$B_\mu B^\mu, \bar{f}\gamma^5f$	$\sigma_{SD} \sim (q/2m_n)^2$	No	Maybe
7	Dirac Fermion	Spin-0 (t-ch.)	$\bar{\chi}(1 \pm \gamma^5)b$	$\sigma_{SI} \sim \text{loop (vector)}$	Yes	Yes
7	Dirac Fermion	Spin-1 (t-ch.)	$\bar{\chi}\gamma^\mu(1 \pm \gamma^5)b$	$\sigma_{SI} \sim \text{loop (vector)}$	Yes	Yes
8	Complex Vector	Spin-1/2 (t-ch.)	$X_\mu^\dagger\gamma^\mu(1 \pm \gamma^5)b$	$\sigma_{SI} \sim \text{loop (vector)}$	Yes	Yes
8	Real Vector	Spin-1/2 (t-ch.)	$X_\mu\gamma^\mu(1 \pm \gamma^5)b$	$\sigma_{SI} \sim \text{loop (vector)}$	Yes	Yes

About half of the tree-level diagrams producing the GC signal are currently compatible with direct detection and collider constraints.

More than 100 papers considering specific models have been submitted.



# FUTURE TESTS OF DARK MATTER

→ DWARF GALAXIES

Dwarf Galaxies can also produce a significant  $\gamma$ -ray signal from dark matter annihilation.

Results from the 4-year, P7 Analysis of the Fermi-LAT data showed a  $TS=8.7$  excess!

FERMI-LAT COLLABORATION (2013, 1310.0828)





# FUTURE TESTS OF DARK MATTER

→ DWARF GALAXIES


However, a new analysis of the Fermi-LAT data was recently presented at the Fermi Symposium (not yet published)

The observed excess has disappeared, and the new limit is now in mild tension with some models of the GC excess



# FUTURE TESTS OF DARK MATTER

NEW DWARFS!




**Analyses of the DES, and Pan-Starrs Data have recently observed 11 (and counting) new dwarf candidates in the Southern Hemisphere.**



# FUTURE TESTS OF DARK MATTER

RETICULUM 2



Several of these dwarfs (Reticulum 2 and possibly Triangulum 2) are close enough to be important targets for dwarf galaxy searches

# FUTURE TESTS OF DARK MATTER



## STELLAR KINEMATICS AND METALLICITIES IN THE ULTRA-FAINT DWARF GALAXY RETICULUM II

J. D. SIMON,<sup>1</sup> A. DRLICA-WAGNER,<sup>2</sup> T. S. LI,<sup>3</sup> B. NORD,<sup>2</sup> M. GEHA,<sup>4</sup> K. BECHTOL,<sup>5</sup> E. BALBINOT,<sup>6,7</sup> E. BUCKLEY-GEER,<sup>2</sup>  
H. LIN,<sup>2</sup> J. MARSHALL,<sup>3</sup> B. SANTIAGO,<sup>8,7</sup> L. STRIGARI,<sup>3</sup> M. WANG,<sup>3</sup> R. H. WECHSLER,<sup>9,10,11</sup> B. YANNY,<sup>2</sup> T. ABBOTT,<sup>12</sup>  
A. H. BAUER,<sup>13</sup> G. M. BERNSTEIN,<sup>14</sup> E. BERTIN,<sup>15,16</sup> D. BROOKS,<sup>17</sup> D. L. BURKE,<sup>10,11</sup> D. CAPOZZI,<sup>18</sup>  
A. CARNERO ROSELL,<sup>7,19</sup> M. CARRASCO KIND,<sup>20,21</sup> C. B. D'ANDREA,<sup>18</sup> L. N. DA COSTA,<sup>7,19</sup> D. L. DEPOY,<sup>3</sup> S. DESAI,<sup>22</sup>  
H. T. DIEHL,<sup>2</sup> S. DODELSON,<sup>2,5</sup> C. E. CUNHA,<sup>10</sup> J. ESTRADA,<sup>2</sup> A. E. EVRARD,<sup>23</sup> A. FAUSTI NETO,<sup>7</sup> E. FERNANDEZ,<sup>24</sup>  
D. A. FINLEY,<sup>2</sup> B. FLAUGHER,<sup>2</sup> J. FRIEMAN,<sup>2,5</sup> E. GAZTANAGA,<sup>13</sup> D. GERDES,<sup>23</sup> D. GRUEN,<sup>25,26</sup> R. A. GRUENDL,<sup>20,21</sup>  
K. HONSCHIED,<sup>27,28</sup> D. JAMES,<sup>12</sup> K. KUEHN,<sup>29</sup> N. KUROPATKIN,<sup>2</sup> O. LAHAV,<sup>17</sup> M. A. G. MAIA,<sup>7,19</sup> M. MARCH,<sup>14</sup>  
P. MARTINI,<sup>27,30</sup> C. J. MILLER,<sup>31,23</sup> R. MIQUEL,<sup>24</sup> R. OGANDO,<sup>7,19</sup> A. K. ROMER,<sup>32</sup> A. ROODMAN,<sup>10,11</sup> E. S. RYKOFF,<sup>10,11</sup>  
M. SAKO,<sup>14</sup> E. SANCHEZ,<sup>33</sup> M. SCHUBNELL,<sup>23</sup> I. SEVILLA,<sup>33,20</sup> R. C. SMITH,<sup>12</sup> M. SOARES-SANTOS,<sup>2</sup> F. SOBREIRA,<sup>2,7</sup>  
E. SUCHYTA,<sup>27,28</sup> M. E. C. SWANSON,<sup>21</sup> G. TARLE,<sup>23</sup> J. THALER,<sup>34</sup> D. TUCKER,<sup>2</sup> V. VIKRAM,<sup>35</sup> A. R. WALKER,<sup>12</sup> AND  
W. WESTER<sup>2</sup>

(THE DES COLLABORATION)

galaxy known. Although Ret II is the third-closest dwarf galaxy to the Milky Way, the line-of-sight integral of the dark matter density squared is  $\log_{10}(J) = 18.8 \pm 0.6 \text{ GeV}^2 \text{ cm}^{-5}$  within  $0.2^\circ$ , indicating that the predicted gamma-ray flux from dark matter annihilation in Ret II is lower than that of several other dwarf galaxies.

**Yeoman's work by several optical spectroscopists has given us two estimations of the J-factors for Reticulum 2**

# FUTURE TESTS OF DARK MATTER



## RETICULUM 2

### DARK MATTER ANNIHILATION AND DECAY PROFILES FOR THE RETICULUM II DWARF SPHEROIDAL GALAXY

VINCENT BONNIVARD<sup>1</sup>, CÉLINE COMBET<sup>1</sup>, DAVID MAURIN<sup>1</sup>, ALEX GERINGER-SAMETH<sup>2</sup>, SAVVAS M. KOUSHIAPPAS<sup>3</sup>,  
MATTHEW G. WALKER<sup>2</sup>, MARIO MATEO<sup>4</sup>, EDWARD W. OLSZEWSKI<sup>5</sup>, AND JOHN I. BAILEY III<sup>4</sup>

*Draft version April 14, 2015*

$\alpha_{\text{int}}$ [deg]	$\log_{10}(J(\alpha_{\text{int}}))$ [ $J/\text{GeV}^2 \text{cm}^{-5}$ ] <sup>a</sup>
0.01	$16.9^{+0.5(+1.1)}_{-0.4(-0.8)}$
0.05	$18.2^{+0.5(+1.0)}_{-0.4(-0.7)}$
0.1	$18.6^{+0.6(+1.1)}_{-0.4(-0.8)}$
0.5	$19.5^{+1.0(+1.6)}_{-0.6(-1.3)}$
1	$19.7^{+1.2(+2.0)}_{-0.9(-1.5)}$

against several of its ingredients. We find that Ret II presents one of the largest annihilation  $J$ -factors among the Milky Way's dSphs, possibly making it one of the best targets to constrain the DM particle properties. However, it is important to obtain follow-up photometric and spectroscopic data in order to test the assumptions of dynamical equilibrium as well as a negligible fraction of binary stars in the kinematic sample. Nevertheless, the proximity of Ret II and its potential large dark matter content make it the most interesting object from the newly discovered dwarf galaxies.


Yeoman's work by several optical spectroscopists has given us two estimations of the  $J$ -factors for Reticulum 2

# FUTURE TESTS OF DARK MATTER

RETICULUM 2

## Search for Gamma-Ray Emission from DES Dwarf Spheroidal Galaxy Candidates with Fermi-LAT Data

A. Drlica-Wagner,<sup>1,2,\*</sup> A. Albert,<sup>3,†</sup> K. Bechtol,<sup>1,4,‡</sup> M. Wood,<sup>3,§</sup> L. Strigari,<sup>5,¶</sup> M. Sánchez-Conde,<sup>6,7</sup> L. Baldini,<sup>8</sup> R. Essig,<sup>9</sup> J. Cohen-Tanugi,<sup>10</sup> B. Anderson,<sup>11</sup> R. Bellazzini,<sup>12</sup> E. D. Bloom,<sup>3</sup> R. Caputo,<sup>13</sup> C. Cecchi,<sup>14,15</sup> E. Charles,<sup>3</sup> J. Chiang,<sup>3</sup> J. Conrad,<sup>7,6,11,16</sup> A. de Angelis,<sup>17</sup> S. Funk,<sup>3</sup> P. Fusco,<sup>18,19</sup> F. Gargano,<sup>19</sup> N. Giglietto,<sup>18,19</sup> F. Giordano,<sup>18,19</sup> S. Guiriec,<sup>20,21</sup> M. Gustafsson,<sup>22</sup> M. Kuss,<sup>12</sup> F. Loparco,<sup>18,19</sup> P. Lubrano,<sup>14,15</sup> N. Mirabal,<sup>20,21</sup> T. Mizuno,<sup>23</sup> A. Morselli,<sup>24</sup> T. Ohsugi,<sup>23</sup> E. Orlando,<sup>3</sup> M. Persic,<sup>25,26</sup> S. Rainò,<sup>18,19</sup> F. Spada,<sup>12</sup> D. J. Suson,<sup>27</sup> G. Zaharijas,<sup>28,29</sup> and S. Zimmer<sup>7,6</sup>  
(The Fermi-LAT Collaboration)



tion 6 in Ackermann *et al.* [19]). The most significant excess for any of the DM masses, annihilation channels, and targets we consider here was  $\text{TS} = 6.7$ , corresponding to a local significance<sup>6</sup> of  $1.5\sigma$  ( $p = 0.06$ ) and a global significance of  $0.26\sigma$  ( $p = 0.40$ ). This coincides with

**Reticulum 2 also has an excess!**

# FUTURE TESTS OF DARK MATTER

## RETICULUM 2


### Evidence for Gamma-ray Emission from the Newly Discovered Dwarf Galaxy Reticulum 2

Alex Geringer-Sameth\* and Matthew G. Walker†  
*McWilliams Center for Cosmology, Department of Physics,  
Carnegie Mellon University, Pittsburgh, PA 15213, USA*

Savvas M. Koushiappas†  
*Department of Physics, Brown University, Providence, RI 02912, USA*

Sergey E. Kopoulos, Vasily Belokurov, Gabriel Torrealba, and N. Wyn Evans  
*Institute of Astronomy, University of Cambridge, Cambridge, CB3 0HA, UK*  
(Dated: March 10, 2015)


We present a search for  $\gamma$ -ray emission from the direction of the newly discovered dwarf galaxy Reticulum 2. Using Fermi-LAT data, we detect a signal that exceeds expected backgrounds between  $\sim 2 - 10$  GeV and is consistent with annihilation of dark matter for particle masses less than a few  $\times 10^2$  GeV. Modeling the background as a Poisson process based on Fermi-LAT diffuse models, and taking into account trials factors, we detect emission with  $p$ -value less than  $9.8 \times 10^{-5}$  ( $> 3.7\sigma$ ). An alternative, model-independent treatment of background reduces the significance, raising the  $p$ -value to  $9.7 \times 10^{-3}$  ( $2.3\sigma$ ). Even in this case, however, Reticulum 2 has the most significant  $\gamma$ -ray signal of any known dwarf galaxy. If Reticulum 2 has a dark matter halo that is similar to those inferred for other nearby dwarfs, the signal is consistent with the  $s$ -wave relic abundance cross section for annihilation.



**Reticulum 2 also has an excess!**

# FUTURE TESTS OF DARK MATTER

RETICULUM 2



Reticulum 2 also has an excess!

# FUTURE TESTS OF DARK MATTER

## DWARF GALAXIES: MODEL BUILDING

If the tension between the GC and dwarf observations persists, this could be addressed via secondary emission models:





The spectrum and morphology of the signal can then be reproduced through the secondary up-scattering of the ISRF.

This is a natural solution in models of self-interacting dark matter.

KAPLINGHAT, TL, YU (2014, 1311.6524)

KAPLINGHAT, TL, YU (2015, 1501.03507)




# FUTURE TESTS OF DARK MATTER



## DIRECT DETECTION SEARCHES

However, these limits are model dependent.

Annihilations through a pseudo-scalar mediator will be unobservable with direct detection






# CONCLUSIONS


- 1.) A bright, spherically symmetric, hard spectrum excess has been observed coincident with the dynamical center of the Milky Way.**
- 2.) This excess is difficult to explain with known astrophysical source mechanisms, such as MSPs and galactic outbursts.**
- 3.) Dark matter provides a natural fit to the characteristics of the GC excess**
- 4.) However, any dark matter claim must be backed up by redundant observations. Significant work must still be done to test out or confirm our models of the GC excess.**

**EXTRA SLIDES**


# COMPARISON TO OTHER RESIDUALS




# FRACTIONAL INTENSITY




# IG EXCESS WITH GC EVENTS REMOVED





# SPECTRUM INSIDE/OUTSIDE BUBBLES



# SPECTRAL VARIATION INSIDE BUBBLES




# CORRELATION WITH GAS






# SPECTRAL VARIATIONS IN DIFFUSE MODEL





# ELLIPTICITY IN GENERAL DIRECTION



# PREVIOUS WORK




GOODENOUGH & HOOPER (2009), HOOPER & GOODENOUGH (2011)



# PREVIOUS WORK

HOOPER & TL (2011)



Employ analytic model for the integrated gas density near the galactic center (Kalberla & Kerp 2009)

Fit the emission in regions far from the galactic center ( $|l| > 5^\circ$ ), and extrapolate into center

Remove emission correlating with gas, and examine intensity and spectrum of remaining emission

# PREVIOUS WORK

→ ABAZAJIAN & KAPLINGHAT (2011)


Produce full model of  $\gamma$ -ray emission in the GC, including all point sources and diffuse emission models

Fit data with, and without, a dark matter component, use log-likelihood to determine best fit

$$\ln \mathcal{L} = \sum_i k_i \ln \mu_i - \mu_i - \ln(k_i!)$$

$\mu_i$  = model counts

$k_i$  = data counts




# PREVIOUS WORK



GORDON & MACIAS (2013), MACIAS & GORDON (2013)



Use Log-Likelihood Formulation, but add additional components corresponding to known high-energy emission sources (20 cm lines, H.E.S.S. ridge)



# PREVIOUS WORK

→ ABAZAJIAN ET AL. (2014)

Examined the variation in the low-energy spectrum of the GC Excess for different choices in the diffuse background modeling.



## The Characterization of the Gamma-Ray Signal from the Central Milky Way: A Compelling Case for Annihilating Dark Matter

Tansu Daylan,<sup>1</sup> Douglas P. Finkbeiner,<sup>1,2</sup> Dan Hooper,<sup>3,4</sup> Tim Linden,<sup>5</sup>  
Stephen K. N. Portillo,<sup>2</sup> Nicholas L. Rodd,<sup>6</sup> and Tracy R. Slatyer<sup>6,7</sup>

<sup>1</sup>*Department of Physics, Harvard University, Cambridge, MA*

<sup>2</sup>*Harvard-Smithsonian Center for Astrophysics, Cambridge, MA*

<sup>3</sup>*Fermi National Accelerator Laboratory, Theoretical Astrophysics Group, Batavia, IL*

<sup>4</sup>*University of Chicago, Department of Astronomy and Astrophysics, Chicago, IL*

<sup>5</sup>*University of Chicago, Kavli Institute for Cosmological Physics, Chicago, IL*

<sup>6</sup>*Center for Theoretical Physics, Massachusetts Institute of Technology, Boston, MA*

<sup>7</sup>*School of Natural Sciences, Institute for Advanced Study, Princeton, NJ*

Past studies have identified a spatially extended excess of  $\sim 1\text{--}3$  GeV gamma rays from the region surrounding the Galactic Center, consistent with the emission expected from annihilating dark matter. We revisit and scrutinize this signal with the intention of further constraining its characteristics and origin. By applying cuts to the *Fermi* event parameter CTBCORE, we suppress the tails of the point spread function and generate high resolution gamma-ray maps, enabling us to more easily separate the various gamma-ray components. Within these maps, we find the GeV excess to be robust and highly statistically significant, with a spectrum, angular distribution, and overall normalization that is in good agreement with that predicted by simple annihilating dark matter models. For example, the signal is very well fit by a 31-40 GeV dark matter particle annihilating to  $b\bar{b}$  with an annihilation cross section of  $\sigma v = (1.4 - 2.0) \times 10^{-26} \text{ cm}^3/\text{s}$  (normalized to a local dark matter density of  $0.3 \text{ GeV}/\text{cm}^3$ ). Furthermore, we confirm that the angular distribution of the excess is approximately spherically symmetric and centered around the dynamical center of the Milky Way (within  $\sim 0.05^\circ$  of Sgr A\*), showing no sign of elongation along or perpendicular to the Galactic Plane. The signal is observed to extend to at least  $\simeq 10^\circ$  from the Galactic Center, disfavoring the possibility that this emission originates from millisecond pulsars.

PACS numbers: 95.85.Pw, 98.70.Rz, 95.35.+d; FERMILAB-PUB-14-032-A, MIT-CTP 4533

### I. INTRODUCTION

Weakly interacting massive particles (WIMPs) are a

tons), other explanations have also been proposed. In particular, it has been argued that if our galaxy's central stellar cluster contains several thousand unresolved mil-



# DAYLAN ET AL. (2014)



## TWO ANALYSIS METHODS

### INNER GALAXY

- Mask galactic plane (e.g.  $|b| > 1^\circ$ )
- Bright point sources masked at  $2^\circ$
- Allow diffuse templates (galactic diffuse, isotropic, Fermi bubbles, dark matter) to float independently in each of 30 energy bins

### GALACTIC CENTER





- Box around the GC ( $10^\circ \times 10^\circ$ )
- Include and model all point sources
- Use likelihood analysis to calculate the spectrum and intensity of each source component
- Calculate log-likelihood to determine significance

# DAYLAN ET AL. (2014)

see Portillo & Finkbeiner (1406.0507)

CTBCORE




Use additional information to classify each photon event based on the accuracy of its directional reconstruction



# PREVIOUS WORK

HOOPER & SLATYER (2013)

Instead analyzed the Fermi bubbles. They found an excess low-energy emission which fell off with increasing distance from the GC.





# FUTURE TESTS OF DARK MATTER

## THE GALACTIC CENTER

Better constraints on the spherical symmetry, spatial extension, and low-energy spectrum of the GC excess can support a DM interpretation.

One interesting analysis has found evidence of a secondary inverse-Compton component with an intensity matching that expected by dark matter annihilation to leptonic final states.




# FERMI-LAT COLLABORATION

Though no Fermi-LAT publication on the GC has yet been published, the preliminary results were shown at 2014 Fermi Symposium.

They also find improved fits when an NFW template is added, the spectral details of the additional component depend on the modeling of the astrophysical diffuse emission.

Simona Murgia, 2014 Fermi Symposium



# FUTURE TESTS OF DARK MATTER



## OTHER GAMMA-RAY TARGETS



May find other bright indirect detection targets.

One possibility is the population of High Velocity Clouds orbiting the Milky Way

Some may be confined by dark matter halos

However, no  $\gamma$ -ray excess is observed in these systems

NICHOLS & BLAND-HAWTHORN (2009, 0911.0684)  
NICHOLS ET AL. (2014, 1404.3209)  
DRLICA-WAGNER ET AL. (2014, 1405.1030)





# FUTURE TESTS OF DARK MATTER

## COSMIC-RAY SEARCHES

Observations of the cosmic-ray positron spectrum by the AMS-02 instrument can place strong constraints on the annihilation to leptonic final states.

In some cases (i.e. direct annihilation to  $e^+e^-$ ) these can fall below the thermal cross-section by two orders of magnitude.

BERGSTROM ET AL. (2013, 1306.3983)





# FUTURE TESTS OF DARK MATTER



## COSMIC-RAY SEARCHES

HOOPER, TL, MERTSCH (2014, 1410.1527)


Observations of Cosmic-Ray Antiproton Fluxes show some evidence for an excess compared to astrophysical models, which can be fit by a dark matter candidate.





# FUTURE TESTS OF DARK MATTER

## DIRECT DETECTION SEARCHES





The 20 - 60 GeV Mass Range is optimal for direct detection searches.

# DAYLAN ET AL. (2014)



## RESULTS: CORRELATION WITH GAS





Can add in the SFD dust map, integrated over the line of sight, and globally bias each ring in order to test the fit to local peaks in the gas density

# CALORE ET AL. (2014) (1409.0042)

Tour de force paper which investigates the resiliency of the  $\gamma$ -ray excess to changes in the astrophysical diffuse model.

Tests over 300 diffuse models and finds the GC excess to be a resilient feature


Finds some evidence for extra high energy emission compared to Daylan et al. (2014)




# FERMI-LAT COLLABORATION

Pulsars, tuned-index


**Without NFW:**



**DATA-MODEL**






Counts in  $0.1^\circ \times 0.1^\circ$  pixels  
 $0.3^\circ$  radius gaussian smoothing



Pulsars, tuned-index

**With NFW:**




# FUTURE TESTS OF DARK MATTER

↪ DWARF GALAXIES

How can this test statistic be translated into a significance?

Can cross-correlate hotspots in the Fermi-LAT data with the positions of known high-energy blazars and radio galaxies.

This allows for a determination of the significance, which was nearly  $2.7\sigma$




# INDIRECT DETECTION OF WIMPS

→ DARK MATTER DENSITY PROFILES

→ THE GALACTIC CENTER: BACKGROUNDS

## What Are These Backgrounds?

- \* Point Sources (SNR, pulsars, etc.)
- \* Hadronic Interactions ( $pp \rightarrow \pi^0 \rightarrow \gamma\gamma$ )
- \* Bremsstrahlung
- \* Inverse Compton Scattering





# PREVIOUS WORK

→ CONSISTENCY!

Despite different background models, ROIs, and degrees of freedom, the results of each analysis are statistically consistent

Hooper & TL (2011)



channel, $m_\chi$	$TS_{\approx}$	$-\ln \mathcal{L}$	$\Delta \ln \mathcal{L}$
$b\bar{b}$ , 10 GeV	2385.7	139913.6	156.5
$b\bar{b}$ , 30 GeV	3460.3	139658.3	411.8
$b\bar{b}$ , 100 GeV	1303.1	139881.1	189.0
$b\bar{b}$ , 300 GeV	229.4	140056.6	13.5
$b\bar{b}$ , 1 TeV	25.5	140108.2	-38.0
$b\bar{b}$ , 2.5 TeV	7.6	140114.2	-44.0
$\tau^+\tau^-$ , 10 GeV	1628.7	139787.7	282.5
$\tau^+\tau^-$ , 30 GeV	232.7	140055.9	14.2
$\tau^+\tau^-$ , 100 GeV	4.10	140113.4	-43.3

Abazajian & Kaplinghat (2012)